

NO-A106 727


OPTIMAL CORRECTION PROBLEM OF A MULTIDIMENSIONAL  
STOCHASTIC SYSTEM(U) WAYNE STATE UNIV DETROIT MI  
J L MENALDI ET AL. SEP 87 AFOSR-TR-87-1479


1/1

UNCLASSIFIED AFOSR-87-0278

F/G 12/3

NL





MICROGRAPH RESOLUTION TEST CHART  
 NATIONAL BUREAU OF STANDARDS-1963-A

AD-A186 727

DTIC FILE COPY

## DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

		1b. RESTRICTIVE MARKINGS	
1. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2. DECLASSIFICATION / DOWNGRADING SCHEDULE			
PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S) <b>AFOSR-TR-87-1479</b>	
1a. NAME OF PERFORMING ORGANIZATION	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION	
2. ADDRESS (City, State, and ZIP Code)		7b. ADDRESS (City, State, and ZIP Code)	
1. NAME OF FUNDING / SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER <b>AFOSR-87-0278</b>	
2. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
1. TITLE (Include Security Classification)			
2. PERSONAL AUTHOR(S)			
1a. TYPE OF REPORT	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day)	15. PAGE COUNT
3. SUPPLEMENTARY NOTATION			
7. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP		
9. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<div style="text-align: center;"><b>DTIC ELECTED</b> OCT 27 1987 KE</div>			
10. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION	
2a. NAME OF RESPONSIBLE INDIVIDUAL		22b. TELEPHONE (Include Area Code)	22c. OFFICE SYMBOL

**AFOSR-TR- 87 - 1 4 7 9**

**OPTIMAL CORRECTION PROBLEM OF  
A MULTIDIMENSIONAL STOCHASTIC SYSTEM**

Approved for public release;  
distribution unlimited.

FOR THE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
OFFICE OF TRANSMISSION AND DTIC  
This technical report has been reviewed and is  
approved for public release (AFR 190-12).  
Distribution is unlimited.  
by **MATTHEW J. KEEPER**  
Chief, Technical Information Division

**J. L. Menaldi\***  
Department of Mathematics  
Wayne State University  
Detroit, MI 48202

and

**M. I. Taksar†**  
Department of Statistics  
Florida State University  
Tallahassee, FL 32306

September, 1987



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

\* Supported by NSF Grant DMS 86-01998.

† Supported by NSF Grant DMS 86-01510 and Air Force Office of Scientific Research Grant 87-0278.

# OPTIMAL CORRECTION PROBLEM OF A MULTIDIMENSIONAL STOCHASTIC SYSTEM

by

J. L. Menaldi\*

Wayne State University

and

M. I. Taksar†

Florida State University

## ABSTRACT

We consider a stochastic dynamic system which is governed by a multidimensional diffusion process with constant drift and diffusion coefficients. The correction corresponds to an additive input which is under control. There is no limit on the rate of input into the system. The objective is to minimize the expected cumulative cost associated with the position of the system and the amount of control exerted.

It is proved that Hamilton-Jacobi-Bellman's equation of the problem has a solution, which corresponds to the optimal cost of the problem. An existence of optimal policy is proved.

---

\* Supported by NSF Grant DMS 86-01998.

† Supported by NSF Grant DMS 86-01510 and Air Force Office of Scientific Research Grant 87-0278.

## 1. INTRODUCTION

This problem is motivated by studies of a dissipative system under uncertainty. A typical model would be an automatic cruise control of an aircraft subject to uncertain wind conditions. The problem is to balance costs associated with deviation of the airplane from the prescribed course and fuel expenditures resulting from the correction of the course.

We assume that in absence of control the fluctuations of our stochastic system are described by a multidimensional Brownian motion with constant  $n$ -dimensional vector - drift  $g$  and  $n \times n$  diffusion matrix  $\sigma$ .

$$y_x(t) = x + gt + \sigma w(t)$$

Here  $x$  is the initial position, and  $w(\cdot)$  is a  $n$ -dimensional standard Brownian motion on  $(\Omega, \mathcal{F}, \mathcal{F}_t, P)$ .

The "quality" of the position of the system is measured by a function  $h$ . We assume that  $h$  is a strictly convex nonnegative function such that

$$h(x)/|x| \rightarrow \infty \text{ as } |x| \rightarrow \infty.$$

The control is realized by  $2n$  increasing,  $\mathcal{F}_t$ -adapted processes  $\nu_i^+(t), \nu_i^-(t), i = 1, 2, \dots, n$ . The control functional  $\nu(t)$  is a  $n$ -dimensional  $\mathcal{F}_t$ -adapted process of bounded variation defined

$$\nu(t) = (\nu_1(t), \nu_2(t), \dots, \nu_n(t)), \quad (1.1)$$

$$\nu_i(t) = \nu_i^+(t) - \nu_i^-(t). \quad (1.2)$$

The dynamics of the system under control is then

$$y_x(t) = x + gt + \sigma w(t) + \nu(t). \quad (1.3)$$

With each initial position  $x$  and each control functional  $\nu$  we associate a cost

$$J_x(\nu) = E \left\{ \int_0^\infty e^{-\alpha t} h(y_x(t)) dt + \sum_{i=1}^n \left[ a_i \int_0^\infty e^{-\alpha t} d\nu_i^+(t) + b_i \int_0^\infty e^{-\alpha t} d\nu_i^-(t) \right] \right\}, \quad (1.4)$$

where  $a_i$  and  $b_i$ ,  $i = 1, 2, \dots, n$  are positive constants and  $\alpha > 0$  is a discount factor. Denote by  $V$  the set of all  $n$ -dimensional  $\mathcal{F}_t$ -adapted processes  $\nu$  represented in the form (1.1), (1.2). We are looking for

$$u(x) = \inf \{ J_x(\nu) : \nu \in V \} \quad (1.5)$$

and  $\nu^*$  such that

$$u(x) = J_x(\nu^*). \quad (1.6)$$

Let

$$[a_{ij}] = \sigma \sigma^* \quad (1.7)$$

and  $\nabla = (\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \dots, \frac{\partial}{\partial x_n})$ . Put

$$A = -\frac{1}{2} \sum_{i,j=1}^n a_{ij} \frac{\partial^2}{\partial x_i \partial x_j} - \sum_{i=1}^n g_i \frac{\partial}{\partial x_i} + \alpha \equiv -\frac{1}{2} \text{tr}(\sigma \sigma^* \nabla^2) - g \nabla + \alpha. \quad (1.8)$$

For  $q = (q_1, q_2, \dots, q_n) \in \mathbb{R}^n$  let  $\|q\| = \max |q_i|$ , Put

$$\beta(q) = \max_{\substack{\xi, \eta \in \mathbb{R}^n; \xi, \eta \geq 0 \\ \|\xi\|, \|\eta\| \leq 1}} \sum_{i=1}^n [-\xi_i(a_i + q_i) - \eta_i(b_i - q_i)] \quad (1.9)$$

Note that

$$\beta(q) \geq 0 \text{ and } \beta(q) = 0 \text{ iff } -a_i \leq q_i \leq b_i, i = 1, 2, \dots, n. \quad (1.10)$$

We will show that the optimal cost function  $u$  given by (1.5) satisfies that following Hamilton-Jacobi-Bellman equation

$$\max(Au - h, \gamma(\nabla u) - 1) = 0, \quad (1.11)$$

where  $\gamma(q) = \max_{1 \leq i \leq n} [(q_i^+ / a_i) \vee (q_i^- / b_i)]$ , and we prove the existence of the optimal cost.

In section 2 we consider a family of problems in which the allowable controls are absolutely continuous with the rates uniformly bounded. We derive estimates for the cost functions of these problems. In section 3 it is shown that a subsequence of cost functions for absolutely continuous control problems converges to  $u(x)$  given by (1.5). Section 4 is devoted to construction of the optimal policy.

## 2. Absolutely continuous control problems.

Here and in sequel we assume that the function  $h$  is strictly convex and roughly speaking is of polynomial growth. More precisely there exist  $p > 1$  and constants  $C_0, C_1$  and  $C_2$  such that for any  $0 < \lambda < 1$ , any  $x \in \mathbb{R}^n$  and any  $x'$  such that  $|x'| \leq 1$ ,

$$0 \leq h(x) \leq C_0(1 + |x|)^p, \quad (2.1)$$

$$|h(x) - h(x + x')| \leq C_1(1 + h(x) + h(x + x'))^{1-p^{-1}} |x'|, \quad (2.2)$$



$$0 < h(x + \lambda x') + h(x - \lambda x') - 2h(x) \leq C_2 \lambda^2 (1 + h(x))^q, \quad q = (1 - \frac{2}{p})^+. \quad (2.3)$$

Let  $V_\epsilon$  be the set of all  $\nu \in V$  such that

$$\nu(t) = \int_0^t \dot{\nu}(s) ds, \quad |\dot{\nu}(s)| < \epsilon^{-1} \quad \text{for all } s > 0, \quad (2.4)$$

and

$$u_\epsilon(x) = \inf_{\nu \in V_\epsilon} J_x(\nu) \quad (2.5)$$

Formal application of the dynamic programming principle yields the following equation

$$Au_\epsilon + \epsilon^{-1} \beta(\nabla u_\epsilon) = h \quad (2.6)$$

(see Fleming and Rishel (1975)). The next theorem establishes the properties of  $u_\epsilon$  which will be used in sequel.

(2.7) **Theorem.** Suppose  $h(x)$  satisfies (2.1) - (2.3). Then there exist  $\hat{C}_0, \hat{C}_1, \hat{C}_2$  independent of  $\epsilon \in (0, 1]$  such that for each  $\lambda \in (0, 1)$  and each  $x'$  with  $|x'| \leq 1$  the function  $u_\epsilon(x)$  satisfies (2.8)-(2.10) below

$$0 \leq u_\epsilon(x) \leq \hat{C}_0(1 + |x|)^p, \quad (2.8)$$

$$|u_\epsilon(x) - u_\epsilon(x + x')| \leq \hat{C}_1(1 + |x| + |x + x'|)^{p-1} |x'|, \quad (2.9)$$

$$0 \leq u_\epsilon(x + \lambda x') + u_\epsilon(x - \lambda x') - 2u_\epsilon(x) \leq \hat{C}_2 \lambda^2 (1 + |x|)^{(p-2)^+}. \quad (2.10)$$

Moreover (2.8)-(2.10) is also true for  $u$ .

**Proof** 1°. Putting  $\nu \equiv 0$ , we get from (2.5)

$$\begin{aligned}
u_\epsilon(x) &\leq J_x(0) = E \left\{ \int_0^\infty e^{-\alpha t} h(x + \sigma w(t) + gt) dt \right\} \\
&\leq E \left\{ \int_0^\infty e^{-\alpha t} C_0 (1 + |x + \sigma w(t) + gt|^p) dt \right\} \\
&\leq C_0 \alpha^{-1} + C_0 E \left\{ \int_0^\infty e^{-\alpha t} |x + \sigma w(t) + gt|^p dt \right\} \\
&\leq C_0 \alpha^{-1} + C_0 \alpha^{-1} c |x|^p,
\end{aligned}$$

where  $c$  is a constant dependent on  $\alpha, \sigma$  and  $g$ . Putting  $\hat{C}_0 = C_0 \alpha^{-1} \max(1, c)$ , we get (2.8).

2°. Consider

$$\begin{aligned}
u_\epsilon(x) - u_\epsilon(x + x') &= \inf_{\nu} \sup_{\nu'} J_x(\nu) - J_{x+x'}(\nu') \\
&\leq \sup_{\nu'} J_{x+x'}(\nu') - J_x(\nu') = \sup_{\nu} J_x(\nu) - J_{x+x'}(\nu).
\end{aligned}$$

Likewise  $u_\epsilon(x + x') - u_\epsilon(x) \leq \sup_{\nu} J_{x+x'}(\nu) - J_x(\nu)$ . Therefore

$$\begin{aligned}
|u_\epsilon(x) - u_\epsilon(x + x')| &\leq \sup_{\nu} |J_x(\nu) - J_{x+x'}(\nu)| \\
&\leq \sup_{\nu} E \left\{ \int_0^\infty e^{-\alpha t} |h(y_x(t)) - h(y_{x+x'}(t))| dt \right\} \\
&\leq \sup_{\nu} C_1 E \left\{ \int_0^\infty |1 + h(y_x(t)) - h(y_{x+x'}(t))|^{1-1/p} \right. \\
&\quad \left. |(y_x(t) - y_{x+x'}(t)) e^{-\alpha t} dt \right\} \\
&\leq C_1 |x'| \alpha^{-1/p} \sup_{\nu} (E \left\{ \int_0^\infty |1 + h(y_x(t)) \right. \\
&\quad \left. + h(y_{x+x'}(t))| e^{-\alpha t} dt \right\})^{1-1/p}.
\end{aligned} \tag{2.11}$$

The last inequality in (2.11) is due Hölder's inequality. By virtue of (2.8), we can consider

only those in (2.11)  $\nu$  for which  $E \int_0^\infty h(y_x(t)) e^{-\alpha t} dt \leq C_0(1 + |x|)^p$ . Therefore, applying Hölder's inequality to the last line of (2.11) once more,

$$|u_\epsilon(x) - u_\epsilon(x + x')| \leq |x'| C_1 \alpha^{-1} (2 + C_0/\alpha) (1 + |x| + |x + x'|)^{p-1},$$

whereas (2.9) follows.

3°. Since  $h$  is a convex function, the function  $J_x(\nu)$  is convex in  $(x, \nu)$ . Because  $V_\epsilon$  is a convex set the function  $u_\epsilon(x)$  is convex as well. Therefore the first part of the inequality (2.10) follows. Put  $x_1 = x + \lambda x'$ ,  $x_2 = x - \lambda x'$

$$\begin{aligned} & u_\epsilon(x + \lambda x') + u_\epsilon(x - \lambda x') - 2u_\epsilon(x) \\ &= \inf_{\nu_1} \inf_{\nu_2} \sup_{\nu} \{J_{x_1}(\nu_1) + J_{x_2}(\nu_2) - 2J_x(\nu)\} \\ &\leq \sup_{\nu} J_{x_1}(\nu) + J_{x_2}(\nu) - 2J_x(\nu) \\ &\leq E \left\{ \int_0^\infty e^{-\alpha t} C_2 \lambda^2 (1 + h(y_x(t)))^q dt \right\}. \end{aligned} \tag{2.12}$$

If  $q = 0$  then (2.12) implies (2.10) in an obvious manner. If  $q > 0$  (that is  $p > 2$ ) then by Hölder's inequality

$$\begin{aligned} & E \left\{ \int_0^\infty e^{-\alpha t} (1 + h(y_x(t)))^q dt \right\} \\ &\leq \left( \int_0^\infty e^{-\alpha t} dt \right)^{2/p} \left( E \left\{ \int_0^\infty e^{-\alpha t} [1 + h(y_x(t))]^{(p-2)/p} dt \right\} \right)^{(p-2)/p} \\ &\leq \alpha^{-2/p} (\alpha^{-q} + \hat{C}_0 (1 + |x|)^p)^{(p-2)/p} \end{aligned} \tag{2.13}$$

The last inequality in (2.13) is due to (2.8). Simple analysis show that (2.13) implies (2.10)

The proof of (2.8)-(2.10) for  $u$  is the same.

(2.14) Theorem. The optimal cost  $u_\epsilon(x)$  is the unique solution of (2.6) under the conditions (2.8)-(2.10).

Proof 1°. For a nondegenerate  $\sigma$  the existence of a solution follows from the classical results (see Fleming and Rishel (1975)). If  $\sigma$  is degenerate then consider  $\sigma_\delta = [\sigma, \delta I]$  which is a  $2n \times n$  matrix. Note that

$$\sigma_\delta \sigma_\delta^* = \sigma \sigma^* + \delta^2 I$$

is nondegenerate for all sufficiently small  $\delta$ . Consider a new optimization problem in which  $w$  in (1.3) is a  $2n$ -dimensional standard Brownian motion while  $\sigma$  is replaced by  $\sigma_\delta$ . Let  $u_{\epsilon, \delta}$  be the corresponding optimal cost given by (2.5). Then  $u_{\epsilon, \delta}$  satisfies

$$-\frac{\delta^2}{2} \nabla u_{\epsilon, \delta} + A u_{\epsilon, \delta} + \epsilon^{-1} \beta(\nabla u_{\epsilon, \delta}) = h \quad (2.14)$$

Repeating step by step the proof of the Theorem (2.7), we can see that (2.8)- (2.10) hold with  $\hat{C}_0, \hat{C}_1, \hat{C}_2$  independent of  $\delta > 0$  for all sufficiently small  $\delta$ . It follows that  $u_{\epsilon, \delta}(x), \nabla u_{\epsilon, \delta}(x)$  and  $A u_{\epsilon, \delta}(x)$  is locally uniformly (in  $\delta$ ) bounded in  $x$ . The latter implies existence of a subsequence  $\delta_k \rightarrow 0$  such that  $|\Delta u_{\epsilon, \delta_k}(x)|$  is locally bounded in  $x$  and

$$\begin{aligned} u_{\epsilon, \delta_k}(x) &\rightarrow u_\epsilon(x), \\ \nabla u_{\epsilon, \delta_k}(x) &\rightarrow \nabla u_\epsilon(x), \end{aligned}$$

locally uniformly in  $x$  and  $A u_{\epsilon, \delta_k} \rightarrow A u_\epsilon$  as a distribution (in Schwartz' sense). Passing to a limit in (2.14), we get the validity of (2.6) for  $u_\epsilon$ .

2°. To prove uniqueness assume that there are two solutions  $u_\epsilon$  and  $v_\epsilon$  of (2.6). Let

$$A_0 = -\frac{1}{2} \sum_{i,j} a_{ij} \frac{\partial^2}{\partial x_i \partial x_j} - \sum_i g_i \frac{\partial}{\partial x_i}.$$

(Recall  $A = A_0 + \alpha I$ ). Then

$$A_0(u_\epsilon - v_\epsilon) = -\alpha(u_\epsilon - v_\epsilon) + \epsilon^{-1}(\beta(\nabla v_\epsilon) - \beta(\nabla u_\epsilon)). \quad (2.15)$$

Let  $W(x) = u_\epsilon(x) - v_\epsilon(x)$  and  $w(x) = W(x)\psi(x, \lambda)$ , where  $\psi(x, \lambda) = (\lambda + |x|^2)^{-p}$  (the value for the constant  $\lambda$  will be chosen later). It follows from (2.8) that  $w(x) \rightarrow 0$  as  $|x| \rightarrow \infty$ . Suppose  $w(x) \not\equiv 0$  and  $w(x) > 0$  for some  $x$ . (If  $w(x) < 0$  then consider  $v_\epsilon - u_\epsilon$ ). Let

$$w(x_0) = \max_{x \in \mathbb{R}^n} w(x) > 0. \quad (2.16)$$

Calculations show that

$$A_0\psi(x, \lambda) + [\text{tr}(\sigma\sigma^* \nabla \psi(x, \lambda) \nabla \psi(x, \lambda)^*)]/\psi(x, \lambda) = \delta(x, \lambda)\psi(x, \lambda), \quad (2.17)$$

where  $\sup_x |\delta(x, \lambda)| \rightarrow 0$  if  $\lambda \rightarrow \infty$ , and

$$\beta(\nabla v_\epsilon) - \beta(\nabla u_\epsilon) = \gamma(x, \epsilon)[\nabla v_\epsilon - \nabla u_\epsilon] \equiv -\gamma(x, \epsilon) \nabla W(x), \quad (2.18)$$

where  $\|\gamma(x, \epsilon)\| \leq 1$ . In view of (2.16),  $\nabla w(x_0) = 0$ . Therefore

$$\nabla W(x_0)\psi(x_0, \lambda) = -W(x_0) \nabla \psi(x_0, \lambda). \quad (2.19)$$

and

$$\nabla W(x_0) = -W(x_0) \nabla \psi(x_0, \lambda)/\psi(x_0, \lambda). \quad (2.20)$$

Since  $\nabla \psi(x_0, \lambda)/\psi(x_0, \lambda) \rightarrow 0$  as  $\lambda \rightarrow \infty$ , we can combine (2.20) and (2.19) to get

$$\beta(\nabla v_\epsilon(x_0)) - \beta(\nabla u_\epsilon(x_0)) = W(x_0)\hat{\delta}(\epsilon, \lambda), \quad (2.21)$$

where  $\sup_{0 < \epsilon \leq 1} |\hat{\delta}(\epsilon, \lambda)| \rightarrow 0$  as  $\lambda \rightarrow \infty$ . Also,

$$A_0 w(x_0) = A_0 W(x_0)\psi(x_0, \lambda) + W(x_0)A_0\psi(x_0, \lambda) - \text{tr}(\sigma\sigma^* \nabla W(x_0) \nabla \psi(x_0, \lambda)). \quad (2.22)$$

Applying to the first term in the right hand side of (2.22) equations (2.16), (2.17) and (2.19), we get

$$\begin{aligned} A_0 w(x_0) &= [-\alpha W(x_0) + \varepsilon^{-1} W(x_0) \hat{\delta}(\varepsilon, \lambda)] \psi(x_0, \lambda) + W(x_0) A_0 \psi(x_0, \lambda) \\ &\quad - \text{tr}(\sigma \sigma^* (-\nabla \psi(x_0, \lambda) W'(x_0) - \psi(x_0, \lambda)) \psi(x_0, \lambda)) \\ &= W(x_0) \psi(x_0, \lambda) (-\alpha + \varepsilon^{-1} \hat{\delta}(\varepsilon, \lambda) + \delta(x_0, \lambda)) \end{aligned}$$

Now choose  $\lambda > 0$  such that  $-\alpha + \varepsilon^{-1} \hat{\delta}(\varepsilon, \lambda) + \delta(x_0, \lambda) < 0$ . In view of (2.16) it follows that this implies  $w(x_0) \leq 0$  which is in contradiction with (2.16).

### 3. Solution of the Hamilton-Jacobi-Bellman equation

Passing to a limit in (2.6), as  $\varepsilon \rightarrow 0$  (and assuming for a moment convergence of  $u_\varepsilon$  to  $u$ , and  $Au_\varepsilon$  to  $u, \nabla u$  and  $Au$  respectively), we get inequalities

$$Au(x) \leq h, \quad (3.1)$$

$$-a_i \leq \nabla u_i(x) \leq b_i. \quad (3.2)$$

Assuming also that at each point  $x$  at least one of (3.1) and (3.2) is tight, we get (1.11). In this section we will show that  $u$  given by (1.5) is actually a solution of (1.11).

(3.3) Theorem. There exist  $c_1, c_2 > 0$  such that

$$c_1|x| \leq u(x) \leq c_2(1 + |x|). \quad (3.4)$$

Proof 1°. For  $x \in \mathbb{R}^n$  put

$$\eta(x) = \begin{cases} x/|x| - x, & \text{if } |x| > 2, \\ 0 & \text{otherwise.} \end{cases}$$

Let

$$\hat{\nu}_x(t) = \eta(x)1_{|x|>2} + \sum_{\tau_k \leq t} \eta(x_{\tau_k}), \quad (3.5)$$

where

$$\tau_0 = 0,$$

$$\tau_k = \inf\{t > \tau_{k-1} : |x + gt + \sigma w(t) + \sum_{n=1}^{k-1} \tau_n| = 2\}.$$

The policy  $\hat{\nu}_x(t)$  acts in the following manner. When the process is outside the ball of radius 2 it is instantaneously moved inside the ball of radius 1. Then there is no action until the process reaches the boundary of the ball of radius 2, at which moment it is moved again into the ball of radius 1 and so on. Let

$$\begin{aligned} U(x) = & E\left\{ \int_0^\infty e^{-\alpha t} h(x + gt + \sigma w(t) + \hat{\nu}_x(t)) dt \right. \\ & + \sum_{k=1}^\infty e^{-\alpha \tau_k} \sum_{i=1}^n \left[ a_i (\hat{\nu}_{xi}(\tau_k) - \hat{\nu}_{xi}(\tau_k^-))^+ + \right. \\ & \left. \left. b_i (\hat{\nu}_{xi}(\tau_k) - \hat{\nu}_{xi}(\tau_k^-))^- \right] \right\}, \quad \text{if } |x| < 2. \end{aligned}$$

Then

$$J_x(\hat{\nu}_x) = \begin{cases} U(x) & \text{if } |x| < 2, \\ U(x/|x|) + \sum_{i=1}^n (a_i(x/|x| - x)_i^+ + b_i(x/|x| - x)_i^-), & \text{if } |x| \geq 2. \end{cases} \quad (3.6)$$

It is easy to see that  $U(x)$  is a continuous function and therefore bounded in  $\{x : |x| < 2\}$ . Thus formula (3.6) implies that for  $\hat{\nu}_x$  given by (3.5),  $J_x(\hat{\nu}_x)/|x|$  is bounded, whereas the second inequality in (3.4) follows.

2°. Let  $a = \min(a_i \wedge b_i, i = 1, 2, \dots, n)$ . Let

$$\mathcal{A}_R = \{|gt + \sigma w(t)| < R \text{ for all } 0 \leq t \leq 1\}$$

Choose  $R$  such that

$$P(\mathcal{A}_R) > \frac{1}{2},$$

and

$$h(x)/|x| > a \text{ for all } |x| > R. \quad (3.7)$$

Let  $|x| > 3R$  and  $\nu$  be any policy.

Put  $B = \{|y_x(t)| \equiv |x + gt + \sigma w(t) + \nu(t)| > |x|/3 \text{ for all } t \leq 1\}$ . Then

$$\begin{aligned} J_x(\nu) &\geq e^{-1} E \left\{ \int_0^1 h(y_x(t)) dt + a \sum_{i=1}^n [\nu_i^+(1) + \nu_i^-(1)] \right\} \geq \\ &e^{-1} E \left\{ \int_0^1 h(y_x(t)) 1_B dt + a \sum_{i=1}^n [\nu_i^+(1) + \nu_i^-(1)] (1 - 1_B) 1_{\mathcal{A}_R} \right\}. \end{aligned} \quad (3.8)$$

We have  $1_B h(y_x(t)) > 1_B a|x|/3$  by virtue of (3.7). On  $\mathcal{A}_R$  the quantity  $|x + gt + \sigma w(t)|$  exceeds  $2|x|/3$  for all  $t \leq 1$ . Therefore on  $\bar{B} \cap \mathcal{A}_R$

$$\sum_{i=1}^n (\nu_i^+(1) + \nu_i^-(1)) > |x|/3.$$

Hence (3.8) exceeds

$$e^{-1} E \left\{ \int_0^1 (a|x|/3) 1_B dt + (a|x|/3) (1 - 1_B) 1_{\mathcal{A}_R} \right\} \geq e^{-1} (a|x|/3) P(\mathcal{A}_R) \geq e^{-1} a|x|/6 \quad (3.9)$$

Inequality (3.9) implies the first inequality in (3.4).

(3.10) **Theorem.** There exists a constant  $c$  and a sequence  $\varepsilon_k \downarrow 0$  such that for any  $R > 0$  there exists  $N$  such that for every  $k > N$



$$u_{\varepsilon_k}(x) < c(1 + |x|) \text{ for all } |x| < R.$$

Proof. Consider a space  $\mathcal{V}$

$$\mathcal{V} = \{v : v \in L^2_\psi, |\nabla v| \in L^2_{\tilde{\psi}}\},$$

where  $\psi = (\lambda + |x|^2)^{-p-n}$ ,  $\tilde{\psi} = (\lambda + |x|^2)^{-p-n+1}$  and  $L^2_\psi$  denote the set of functions  $v$  on  $\mathbb{R}^n$  for which  $v^2\psi$  is integrable.

Inequalities (2.8)-(2.10) show that  $u_\varepsilon(x)$ ,  $\nabla u_\varepsilon(x)$  and  $\|\frac{\partial^2 u_\varepsilon(x)}{\partial x^2}\|$  are uniformly bounded in  $\varepsilon$  on every compact subset of  $\mathbb{R}^n$ . Also the same inequalities show that  $u_\varepsilon$  is uniformly bounded in  $L^2_\psi$ ,  $|\nabla u_\varepsilon|$  is uniformly bounded in  $L^2_{\tilde{\psi}}$  and  $\|\frac{\partial^2 u_\varepsilon(x)}{\partial x^2}\|$  is uniformly bounded in  $L^2_\psi$ . Hence there exists a function  $u_0 \in \mathcal{V}$  and a subsequence  $\varepsilon_k$  such that

$$u_{\varepsilon_k} \rightarrow u_0 \text{ weakly in } \mathcal{V}, \quad (3.11)$$

$$Au_{\varepsilon_k} \rightarrow Au_0 \text{ weakly in } L^2_\psi \quad (3.12)$$

and  $\varepsilon_k^{-1}\beta(\nabla u_{\varepsilon_k}) = h - Au_{\varepsilon_k}$  is bounded in  $L^2_\psi$ . Therefore

$$\lim_{k \rightarrow \infty} \beta(\nabla u_{\varepsilon_k}) = 0. \quad (3.13)$$

Since convergence of  $\nabla u_{\varepsilon_k}$  is locally uniform, by virtue of (1.10), for every  $R$  there exists  $N$  such that for every  $k > N$

$$-a_i - \delta < \nabla u_{\varepsilon_k i}(x) < b_i + \delta \text{ for all } |x| < R. \quad (3.14)$$

Since  $u_{\varepsilon_k}(x)$  is monotonically decreasing, (3.14) implies the statement of the theorem.

(3.15) Proposition. Let  $V_0$  denote the set of  $\nu \in V$  such that  $\nu(0) = 0$ . then

$$u(x) = \inf_{\nu \in V_0} J_x(\nu).$$

Proof 1°. We may consider only those  $\nu$  for which  $J_x(\nu)$  is finite. First show that in the minimization problem (1.5) we can consider only those  $\nu$  for which there exists  $r$  (possibly dependent on  $x$ ) such that

$$|\nu(0)| < r \text{ a.s.} \quad (3.16)$$

Let  $\hat{\nu}_x$  be a policy given by (3.5) for which  $J_x(\hat{\nu}_x) < c_2(1 + |x|)$ . For any policy  $\nu$  and initial state  $x$  consider a policy

$$\nu^r(t) = \begin{cases} \nu(t) & \text{if } |\nu(0)| < r, \\ \hat{\nu}_x(t) & \text{if } |\nu(0)| \geq r. \end{cases}$$

If  $|r| > |x|$  then  $|\nu^r(0)| < r$ . Using the first inequality in (3.4),

$$\begin{aligned} J_x(\nu) - J_x(\nu^r) &\geq E\{c_1|y_x(0)| - c_2(1 + |x|); |\nu(0)| \geq r\} \\ &\geq (c_1(r - |x|) - c_2(1 + |x|))P(|\nu(0)| \geq r). \end{aligned} \quad (3.17)$$

If  $r > (1 + |x|)c_2/c_1 + |x|$ , then  $J_x(\nu^r) < J_x(\nu)$  while  $|\nu^r(0)| < r$ .

Likewise, we can show that every policy is dominated by the one for which for every stopping time  $\tau$

$$|\nu(y_x(\tau)) - \nu(y_x(\tau-))| \leq (1 + |y_x(\tau-)|)c_2/c_1 + |y_x(\tau-)|. \quad (3.18)$$

2°. Let  $\nu$  be any policy subject to (3.16) and (3.18) and let

$$\tilde{\nu}(t; \epsilon) = \begin{cases} \nu(t) & \text{if } t \geq \epsilon, \\ \nu(0)t/\epsilon + \nu(t)(t - \epsilon)/\epsilon & \text{if } t < \epsilon. \end{cases}$$

Let  $\tilde{y}_x$  be the trajectory corresponding to  $\tilde{\nu}(t; \delta)$  and

$$\tau = \inf\{t : |\nu(t)| > r + 1 \text{ or } |\tilde{y}_x(t)| > |x| + r + 1\}$$

$$\nu(t; \varepsilon) = \begin{cases} \tilde{\nu}(t; \varepsilon), & \text{if } \tau > \varepsilon, \text{ or if } \tau < \varepsilon \text{ and } t \leq \tau, \\ \hat{\nu}_{\tilde{y}_x(\tau)}(t - \tau) + \tilde{\nu}(\tau; \varepsilon), & \text{if } \tau \leq \varepsilon \text{ and } t > \tau, \end{cases}$$

where  $\hat{\nu}$  is given by (3.5). Let  $y_x(t)$  be the trajectory corresponding to  $\nu$  and  $y_x(t; \varepsilon)$  be the trajectory corresponding to  $\nu(\cdot; \varepsilon)$ . Then,  $\nu(t, \varepsilon) = \nu(t)$  for all  $t > \varepsilon$  on  $\{\tau > \varepsilon\}$ . Also, in view of (3.16)

$$P\{\tau < \varepsilon\} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0. \quad (3.19)$$

Consider

$$\begin{aligned} |J_x(\nu) - J_x(\nu(\cdot; \varepsilon))| &\leq E\left\{\int_0^\varepsilon h(y_x(t))e^{-\alpha t}dt; \tau > \varepsilon\right\} \\ &+ E\left\{\int_0^\varepsilon h(y_x(t; \varepsilon))e^{-\alpha t}dt; \tau > \varepsilon\right\} + E\left\{\int_0^\infty h(y_x(t))e^{-\alpha t}dt\right. \\ &+ \sum_{i=1}^n \left[ a_i \int_0^\infty e^{-\alpha t} d\nu_i^+(t) + b_i \int_0^\infty e^{-\alpha t} d\nu_i^-(t) \right]; \tau \leq \varepsilon \Big\} + E\left\{\int_0^\infty h(y_x(t; \varepsilon))e^{-\alpha t}dt\right. \\ &+ \sum_{i=1}^n \left[ a_i \int_0^\infty e^{-\alpha t} d\nu_i^+(t; \varepsilon) + b_i \int_0^\infty e^{-\alpha t} d\nu_i^-(t; \varepsilon) \right]; \tau \leq \varepsilon \Big\} \\ &+ |E\left\{\sum_{i=1}^n a_i \left( \int_0^\varepsilon e^{-\alpha t} d\nu_i^+(t) - \int_0^\varepsilon e^{-\alpha t} d\nu_i^+(t; \varepsilon) \right); \tau > \varepsilon\right\}| \\ &+ |E\left\{\sum_{i=1}^n b_i \left( \int_0^\varepsilon e^{-\alpha t} d\nu_i^-(t) - \int_0^\varepsilon e^{-\alpha t} d\nu_i^-(t; \varepsilon) \right); \tau > \varepsilon\right\}| \\ &= I_1 + I_2 + I_3 + I_4 + I_5 + I_6. \end{aligned} \quad (3.20)$$

The term under expectation in  $I_1$  is majorized by  $\int_0^\infty e^{-\alpha t} h(y_x(t)) dt$  which has finite mean by assumption. Thus  $I_1 \rightarrow 0$  as  $\varepsilon \rightarrow 0$  by virtue of the dominated convergence theorem. Since  $y_x(t; \varepsilon) = \tilde{y}_x(t)$  on  $\{t < \tau\}$ , we have

$$|y_x(t; \varepsilon)| = |\tilde{y}_x(t)| \leq |x| + r + 1 \quad \text{on } \{t \leq \varepsilon < \tau\}.$$

Therefore  $I_2$  does not exceed  $E\{M\varepsilon; \tau > \varepsilon\} \leq M\varepsilon$  where  $M = \max_{|y| \leq |x| + r + 1} h(y)$ .

By assumption  $J_x(\nu) < \infty$  therefore, by virtue of (3.19) and (3.16) and the dominated convergence theorem,  $I_3 \rightarrow 0$ . By virtue of (3.18)

$$|y_x(\tau; \varepsilon)| \leq |x| + r + 1 + (1 + |x| + r + 1)c_2/c_1 \quad (3.21)$$

In view of (3.6) and the strong Markov property for  $y_x(\cdot; \varepsilon)$ , we get that  $I_4$  does not exceed  $E\{M\varepsilon + c_2(1 + |y_x(\tau; \varepsilon)|); \tau \leq \varepsilon\}$ . Therefore (3.19) and (3.21) imply  $I_4 \rightarrow 0$  as  $\varepsilon \rightarrow 0$ . On the set  $\{\tau > \varepsilon\}$  the functional  $\nu_i^+(t), t < \varepsilon$  is bounded by  $|x| + r + 1$ . Likewise for  $\nu_i^-(t; \varepsilon)$ . Straightforward verification shows that both  $\int_0^\varepsilon e^{-\alpha t} d\nu_i^+(t)$  and  $\int_0^\varepsilon e^{-\alpha t} d\nu_i^-(t; \varepsilon)$  converge to  $\nu(0)$  as  $\varepsilon \rightarrow 0$ . Therefore by the bounded convergence theorem  $I_5 \rightarrow 0$  as  $\varepsilon \rightarrow 0$ . Similarly for  $I_6$ .

Let  $V' = \bigcup_{\varepsilon > 0} V_\varepsilon$ .

(3.22) Theorem The set  $V'$  is dense in  $V_0$  that is

$$\inf_{\nu \in V_0} J_x(\nu) = \inf_{\nu \in V'} J_x(\nu).$$

Proof 1°. Let  $\nu \in V$  such that  $J_x(\nu) < \infty$ .

Put

$$\nu_i^\pm(t, \delta) = \begin{cases} 0, & \text{if } 0 \leq t \leq \delta, \\ \delta^{-1} \int_{t-\delta}^t \min(\nu_i^\pm(s), \delta^{-1}) ds, & \text{if } t \geq \delta. \end{cases}$$

It is obvious that  $\nu(t; \delta) \equiv (\nu_1^+(t, \delta) - \nu_1^-(t, \delta), \dots, \nu_n^+(t, \delta) - \nu_n^-(t, \delta))$  is a continuous functional with

$$|\dot{\nu}(t, \delta)| \leq n\delta^{-2}.$$

It is also clear that

$$\nu_i^\pm(t; \delta) \leq \nu_i^\pm(t) \quad (3.23)$$

and  $\nu(t; \delta) \rightarrow \nu(t)$  as  $\delta \rightarrow 0$  for all  $t$  except possibly a countable set of the points of discontinuity of  $\nu$ . To justify convergence of  $J_x(\nu(\cdot, \delta))$  to  $J_x(\nu)$  we, however, need several extra steps as well as a further modification of  $\nu(\cdot, \delta)$ .

Proof 2°. Let  $y_x$  be the trajectory associated with  $\nu$  and  $y_x^\delta$  be the trajectory associated with  $\nu(\cdot, \delta)$  and let

$$\tau_R = \inf\{t : |y_x(t)| \geq R\},$$

$$\tau_R^\delta = \inf\{t : |y_x^\delta(t)| \geq R\}.$$

Fix  $R$ . By virtue of the Theorem (3.10) there exist  $\varepsilon > 0$  and  $\eta_x(t) \in V_\varepsilon$  such that

$$J_x(\eta_x) < c(1 + |x|) \quad \text{for all } |x| < 4nR. \quad (3.24)$$

For every  $\delta$  less than  $\varepsilon$  above, put

$$\sigma(R) = \tau_R \wedge \tau_{4nR}^\delta \wedge \inf\{t : \max_i \nu_i^+(t) \vee \max_i \nu_i^-(t) = N\}$$

(the constant  $N$  will be chosen later) and put

$$\bar{\nu}(t, \delta) = \begin{cases} \nu(t, \delta), & \text{if } t < \sigma(R), \\ \nu(\sigma(R), \delta) + \eta_{y_{\sigma(R)}}^\delta(t - \sigma(R)), & \text{if } t \geq \sigma(R). \end{cases}$$

The policy  $\nu(t, \delta)$  coincides with policy  $\nu(t, \delta)$  which approximates the original policy  $\nu$  until either the process  $|y_x(t)|$  reaches  $R$  or the process  $|y_x^\delta(t)|$  reaches the level  $4nR$  or one of the control functionals  $\nu_i^\pm(\cdot)$  exceeds  $N$ . After that  $\nu(\cdot, \delta)$  switches to the policy whose expected cost grows with  $x$  at the rate not exceeding  $|x|$ . It is obvious that  $\nu(\cdot, \delta) \in V_\delta$ . In the sections 3° and 4° we will show

$$\overline{\lim}_{\substack{\delta \rightarrow 0 \\ R \rightarrow \infty}} J_x(\nu(\cdot, \delta)) \leq J_x(\nu). \quad (3.25)$$

3°. From the definition of  $\nu(\cdot, \delta)$  it is clear that

$$\nu_i^\pm(t-) \leq \overline{\lim}_{\delta \rightarrow 0} \nu_i^\pm(t, \delta) \leq \nu_i^\pm(t).$$

Similarly if  $t_n \rightarrow t$

$$\nu_i^\pm(t-) \leq \overline{\lim}_{\delta \rightarrow 0, t_n \rightarrow t} \nu_i^\pm(t_n, \delta) \leq \nu_i^\pm(t). \quad (3.26)$$

Suppose that with positive probability

$$\lim_{\delta \rightarrow 0} \tau_{4nR}^\delta < \tau_R. \quad (3.27)$$

Let  $w(t)$  be the trajectory for which (3.27) holds. Then there exists a sequence  $\delta_n \downarrow 0$  and a bounded sequence  $t_n < \tau_R$  such that

$$|y_{xi}^{\delta_n}(t_n)| \geq 4R. \quad (3.28)$$

By choosing a subsequence if necessary, we may assume  $t_n \rightarrow t$ . Since  $|y_{xi}(t_n)| < R$  on  $\{t_n < \tau_R\}$ , (3.28) implies

$$\lim_{n \rightarrow \infty} |y_{xi}^{\delta_n}(t_n) - y_{xi}(t_n)| = \lim_{n \rightarrow \infty} |\nu_i(t_n, \delta_n) - \nu_i(t_n)| \geq 3R. \quad (3.29)$$

However  $\nu_i$  is continuous from the right and has left limits. Therefore

$$\nu_i^\pm(t-) \leq \overline{\lim}_{t_n \rightarrow t} \nu_i^\pm(t_n) \leq \nu_i^\pm(t). \quad (3.30)$$

Inequality (3.30) and (3.26) show that

$$\lim |\nu_i(t_n, \delta_n) - \nu_i(t_n)| \leq |\nu_i(t) - \nu_i(t-)| \quad (3.31)$$

however,  $|\nu_i(t) - \nu_i(t-)| = |y_{xi}(t) - y_{xi}(t-)| \leq 2R$  on the set  $\{t < \tau_R\}$ . The latter contradicts to (3.31) and (3.29). Therefore the probability of (3.27) is null.

4°. For any policy  $\nu$  put

$$j_x(t, \nu) = \int_0^t e^{-\alpha t} h(y_x(t)) dt + \sum_{i=1}^n \left[ a_i \int_0^t e^{-\alpha t} d\nu_i^+(t) + b_i \int_0^t e^{-\alpha t} d\nu_i^-(t) \right].$$

Since  $\tau_R \uparrow \infty$  as  $R \rightarrow \infty$ , we can apply the dominated convergence theorem to obtain

$$J_x(\nu) = \lim_{R \rightarrow \infty} E\{j_x(\tau_R, \nu)\},$$

which implies

$$\lim_{R \rightarrow \infty} E\{j_x(\infty, \nu) - j_x(\tau_R, \nu)\} = 0.$$

Applying the strong Markov property for  $y_x(\cdot)$  and the first part of the inequality (3.4), we get

$$E\{e^{-\tau_R} c R\} \leq E\{e^{-\tau_R} c | y_x(\tau_R)\} \rightarrow 0 \quad \text{as} \quad R \rightarrow \infty. \quad (3.32)$$

Consider

$$\begin{aligned}
J_x(\nu(\cdot, \delta)) &= E\{j_x(\infty, \nu(\cdot, \delta))\} \\
&= E\{j_x(\sigma(R), \nu(\cdot, \delta))\} + E\{j_x(\infty, \nu(\cdot, \delta)) - j_x(\sigma(R), \nu(\cdot, \delta))\}
\end{aligned} \tag{3.33}$$

In view of (3.32) there exists  $R$  such that

$$c(1 + 4nR)E\{e^{-\tau_R}\} < \epsilon. \tag{3.34}$$

Then choose  $N$  such that

$$c(1 + 4nR)P\{\max_i \nu_i^+(\tau_R) \vee \max_i \nu_i^-(\tau_R) > N\} < \epsilon. \tag{3.35}$$

since (3.27) does not hold a.s.

$$\sigma(R) \rightarrow \tau_R \wedge \inf\{t : \max_i \nu_i^+(t) \vee \max_i \nu_i^-(t) = N\} \equiv \zeta, \text{ as } \delta \rightarrow 0.$$

Since  $\nu(t, \delta) \rightarrow \nu(t)$  for all  $t < \zeta$  except a countable number of  $t$

$$j_x(\sigma(R), \nu(\cdot, \delta)) \rightarrow j_x(\zeta, \nu). \tag{3.36}$$

Moreover the convergence in (3.36) is bounded because  $|y_x(t)| < R$  and  $|y_x^\delta(t)| < 4R$  and  $\nu_i^\pm(t; \delta) \leq \nu_i^\pm(t) \leq N$  if  $t < \sigma(R)$ . Therefore

$$\lim_{\delta \rightarrow 0} E\{j_x(\sigma(R), \nu(\cdot, \delta))\} \rightarrow E\{j_x(\zeta, \nu)\} \leq J_x(\nu). \tag{3.37}$$

In view of (3.24) and the strong Markov property for  $y_x^\delta$  the second term in (3.33) does not exceed



$$\begin{aligned}
& E\{e^{-\alpha\sigma(R)}c(1 + |y_x^\delta(\sigma(R))|)\} \\
& \leq c(1 + 4R)E\{e^{-\alpha\sigma(R)}\} \\
& \leq c(1 + 4R)\left[E\{e^{-\alpha\tau_R}; \tau_R = \sigma(R)\} \right. \\
& \quad \left. + E\{e^{-\alpha\sigma(R)}; \tau_R > \tau_{4R}^\delta\} \right. \\
& \quad \left. + E\{e^{-\alpha\sigma(R)}; \tau_R > \zeta\}\right] \\
& \leq c(1 + 4R)E\{e^{-\alpha\tau_R}\} + c(1 + 4R)P\{\tau_R > \tau_{4R}^\delta\} \\
& \quad + c(1 + 4R)P\{\tau_R > \zeta\}.
\end{aligned} \tag{3.38}$$

The first term in (3.38) does not exceed  $\varepsilon$  by virtue of (3.34). Since (3.27) holds with probability zero the second term in (3.38) can be made smaller than  $\varepsilon$  when  $\delta$  is sufficiently small. The third term in (3.38) does not exceed  $\varepsilon$  in view of (3.35). In view of arbitrariness of  $\varepsilon$ , we get (3.25). The statement of the theorem is a trivial consequence of (3.25).

(3.39) Corollary. For every  $x$

$$\lim_{\varepsilon \rightarrow 0} u_\varepsilon(x) = u(x).$$

The proof of this theorem follows from Proposition (3.15) and Theorems (3.22).

(3.40) Theorem. The optimal cost  $u$  given by (1.5) satisfies (1.11).

*Proof.* The proof of Theorem (3.10) shows that there exists a function  $u_0$  for which (3.11) and (3.12) hold.

Since  $\beta(q)$  is a continuous function of  $q$  and  $\nabla u_{\varepsilon_k} \rightarrow \nabla u$  we have

$$\beta(\nabla u_0) = 0 \tag{3.41}$$

and in view of (1.10) we get (3.2). From (2.6)

$$Au_\epsilon \leq au_\epsilon + \epsilon^{-1}\beta(\nabla u_\epsilon) = h$$

and it follows (after passing to a limit as  $\epsilon_k \rightarrow 0$ )

$$Au_0 \leq h$$

Suppose  $\gamma(\nabla u_0(x_0)) < 1$ , that is

$$-a_i < \nabla u_0(x_0)_i < b_i \quad \text{for all } 1 \leq i \leq n. \quad (3.42)$$

By virtue of the continuity of  $\nabla u_0$ , the inequality (3.42) is true for all  $|x - x_0| < \delta$  for some  $\delta > 0$  since convergence of  $\nabla u_{\epsilon_k}$  to  $\nabla u_0$  is locally uniform (see the proof of Theorem (3.10)) the inequality (3.42) is true for  $\nabla u_{\epsilon_k}$  for all  $|x - x_0| < \delta$  and all  $k$  sufficiently large. For such  $k$

$$Au_{\epsilon_k} = h$$

and, passing to a limit as  $k \rightarrow \infty$ ,

$$Au_0 = h, \quad (3.43)$$

that is  $\gamma(\nabla u) < 1$  implies (3.43). In view of corollary (3.39),  $u = \lim u_\epsilon$ , hence  $u_0 = u$  and the theorem is proved.

#### 4. Construction of the optimal policy

Let

$$\Delta_2 h(x, y) = (h(x) + h(y))/2 - h((x + y)/2). \quad (4.1)$$

By virtue of (2.3)

$$\Delta_2 h(x, y) > 0, \quad \text{if } x \neq y. \quad (4.2)$$

(4.3) Theorem. The optimal policy  $\nu^*$  (if exists) is unique.

Proof. Suppose there are  $\nu_1^*$  and  $\nu_2^*$  for which (1.6) is true. Let  $y_x^1(t)$  and  $y_x^2(t)$  be the corresponding trajectories. Put  $\nu = (\nu_1^* + \nu_2^*)/2$  and  $y_x(t) = (y_x^1(t) + y_x^2(t))/2$ . Then

$$\begin{aligned} u(x) - J_x(\nu) &= (J_x(\nu_1^*) + J_x(\nu_2^*)) / 2 - J_x(\nu) \\ &\geq E \left\{ \int_0^\infty e^{-\alpha t} \Delta_2 h(y_x^1(t), y_x^2(t)) dt \right\}. \end{aligned} \quad (4.4)$$

By virtue of (4.2) the right hand side of (4.4) is strictly positive if  $\nu_1^*$  and  $\nu_2^*$  are not equal a.s.

Let  $m_T$  be a measure on  $([0, T] \times \Omega, \mathcal{B}[0, T] \times \mathcal{F})$  equal to the product of Lebesgue measure and  $P$ .

(4.5) Theorem. If

$$J_x(\nu_k) \rightarrow u(x) \quad \text{as } k \rightarrow \infty, \quad (4.6)$$

then  $\nu_k(t, \omega)$  converges in measure  $m_T$ .

Proof. 1°. Let  $y_x^k$  be the trajectory corresponding to  $\nu_k$ . then

$$E \left\{ \int_0^T 1_{|y_x^k(t)| > N} dt \right\} \rightarrow 0 \quad \text{as } N \rightarrow \infty \quad (4.7)$$

uniformly in  $k$ . Really

$$\begin{aligned}
J_x(\nu_k) &\geq e^{-\alpha T} E \left\{ \int_0^T h(y_x^k(t)) dt \right\} \\
&\geq e^{-\alpha T} \inf_{|x| \geq N} h(x) E \left\{ \int_0^T 1_{|y_x^k(t)| > N} dt \right\}.
\end{aligned} \tag{4.8}$$

Because  $J_x(\nu_k)$  is uniformly bounded in  $k$  and  $\lim_{N \rightarrow \infty} \inf_{|x| \geq N} h(x) = \infty$ , we get (4.7).

2°. We need to show that for any  $\varepsilon > 0$

$$E \left\{ \int_0^T 1_{|y_x^k(t) - y_x^m(t)| > \varepsilon} dt \right\} \rightarrow 0 \quad \text{as } m, k \rightarrow \infty. \tag{4.9}$$

Suppose that the expectation in (4.9) is greater than  $\delta > 0$  for all  $m$  and  $k$  (or for all  $m$  and  $k$  from a subsequence). Let  $N$  be such that the expectation in (4.7) is less than  $\delta/2$  for all  $k$ . Then

$$\begin{aligned}
&(J_x(\nu_k) + J_x(\nu_m))/2 - J_x((\nu_k + \nu_m)/2) \\
&\geq e^{-\alpha T} E \left\{ \int_0^T \Delta_2 h(y_x^m(t), y_x^k(t)) 1_{|y_x^k(t)| \leq N} 1_{|y_x^m(t)| \leq N} dt \right\} \\
&\geq e^{-\alpha T} \rho(\varepsilon, N) E \left\{ \int_0^T 1_{|y_x^k(t) - y_x^m(t)| > \varepsilon} 1_{|y_x^m(t)| \leq N} 1_{|y_x^k(t)| \leq N} dt \right\} \\
&e^{-\alpha T} \rho(\varepsilon, N) \delta/2,
\end{aligned} \tag{4.10}$$

where  $\rho(\varepsilon, N) = \inf_{|x-y| \geq \varepsilon; |x|, |y| \leq N} \Delta_2 h(x, y)$ . (Usual continuity/compactness arguments show that  $\rho(\varepsilon, N) > 0$  for any  $N$ ). Inequality (4.10) and (4.6) imply  $\lim J_x((\nu_k + \nu_m)/2) < u(x)$  and we come to a contradiction.

(4.11) Corollary. There exists an optimal policy  $\nu^*$ .

Proof Taking a sequence  $\nu_k$  for which (4.5) is true, and using a diagonal method, we can find a subsequence  $\nu_{k_m}$  which converges a.e.  $m_T$  for each  $T > 0$ . Usual argument show an existence of  $\nu$  such that

$$\nu(t, \omega) = \lim \nu_{k_m}(t, \omega) \quad (4.12)$$

for  $\text{Leb} \times P$  almost all  $(t, \omega)$ . By Fatou's lemma

$$\lim J_x(\nu_{k_m}) \geq J_x(\lim(\nu_{k_m})) = J_x(\nu).$$

Thus  $J_x(\nu) \leq u(x)$ . Therefore  $\nu$  given by (4.12) coincides with  $\nu^*$ .

## REFERENCES

- Bather, J.A. and H. Chernoff (1967). Sequential decisions in the control of spaceship. In *Proc. Fifth Berkeley Symposium of Mathematical Statistics and Probability*, Vol. 3, Berkeley, University of California Press, 181-207.
- Bather, J. A. and H. Chernoff (1967). Sequential decisions in the control of a spaceship (finite fuel). *J. Appl. Prob.*, 4, 584-604.
- Benes, V. E., L. A. Shepp and H. S. Witsenhausen (1980). Some solvable stochastic control problems. *Stochastics*, 4, 39-83.
- Bensoussan, A. and J. L. Lions (1978). *Applications des inequations variationnelles en controle stochastique*. Dunod, Paris.
- Billingsley, P. (1968). *Convergence of probability measures*. Wiley.
- Chernoff, H. (1968). Optimal stochastic control. *Sankhyā. Ser. A*, 30, 221-252.
- Chernoff, H. and A. J. Petkau (1978). Optimal control of a Brownian motion. *SIAM J. Appl. Math.* 34, 717-731.
- Chernousko, F. L. (1968). Optimum correction under active disturbances. *Appl. Math. Mech.* (PMM) 32, 203-208.
- Chow, P. L., J. L. Menaldi, M. Robin (1985). Additive control of stochastic linear systems with finite horizon. *SIAM J. on Control and Optimization*, 6, 858-899.
- Fleming, W. H. and R. W. Rishel (1975). *Deterministic and stochastic optimal control*. Springer-Verlag, New York.
- Harrison J. M. and M. I. Taksar (1983). Instantaneous control of a Brownian motion. *Math Oper. Res.*, 8, 439-453.
- Karatzas, I. (1981). The monotone follower problem in stochastic decision theory. *Appl. Math. Optim*, 7, 175-189.
- Karatzas, I. (1983). A class of singular stochastic control problems. *Adv. Appl. Prob.*, 15, 225-254.
- Karatzas, I. and S. E. Shreve (1984). Connection between optimal stopping and singular stochastic control I. Monotone follower problems. *SIAM J. on Control and Optimization*, 22, 856-877.
- Karatzas, I. and S. E. Shreve (1985). Connection between optimal stopping and singular

- stochastic control II. Reflected follower problems. *SIAM J. on Control and Optimization*, **23**, 433-541.
- Lions, P. L. and J. L. Menaldi (1982). Optimal control of stochastic integrals and Hamilton-Jacobi-Bellman equations I. *SIAM J. on Control and Optimization*, **20**, 58-81, 82-95.
- Menaldi, J. L. (1980). On the optimal impulse control problem for degenerate diffusions. *SIAM J. on Control and Optimization*, **18**, 722-739. See also C. R. Acad. Sc. Paris, Serie A, 284 (1977), 1499-1502.
- Menaldi, J. L. and M. Robin (1983). On some cheap control problems for diffusion processes. *Trans. Am. Math. Soc.*, **278**, 771-802.
- Manaldi, J. L. and M. Robin (1984). On singular control problems for diffusions with jumps. *IEEE Trans. Automatic Control*, **AC-29**, 991- 1004.
- Manaldi, J. L. and M. Robin (1985). On optimal correction problems with partial information. *Stoch. Anal. Appl.*, **3**, 63-92.
- Menaldi, J. L. and Taksar M. I. (1987). Singular control of multidimensional Brownian motion, *Proceedings of the 10th IFAC Congress*, Munich, Germany, 1987.
- Meyer, P. A. (1976). *Cours sur les integrales stochastiques*. Lecture Notes in Mathematics 511. Springer-Verlag, new York, 245-400.
- Robin, M. (1978). *Contrôle impulsif des processus de Markov*. These d'Etat, INRIA, Le Chesnay, France. See also C. R. Acad. Sc. Paris, Serie A, 282 (1976), 631-642.
- Taksar, M. I. (1984). Storage model with discontinuous holding cost. *Stoch. proc. Appl.*, **18**, 201-300.
- Taksar, M. I. (1985). Average optimal control and a related optimal stopping problem. *Math. Oper. Res.*, **10**, 63-81.
- Taksar, M. I. (1986). Free boundary control and a related optimal stopping problem. In the *Proceedings of the 25th IEEE Conference on Decision and Control*, Athens, Greece, 1986, 132-133.

END

JAN.

1988

DTIC